

Print ISSN : 0972-8813
e-ISSN : 2582-2780

[Vol. 24(1) January-April 2026]

Pantnagar Journal of Research

(Formerly International Journal of Basic and
Applied Agricultural Research ISSN : 2349-8765)



G.B. Pant University of Agriculture & Technology
Pantnagar, U.S. Nagar; Uttarakhand, Website : gbpuat.res.in/PJR



ADVISORY BOARD

Patron

Prof. Shivendra Kumar Kashyap, Ph.D., Vice-Chancellor, G.B. Pant University of Agriculture and Technology, Pantnagar, India

Members

Prof. S. K. Verma, Ph.D., Director Research, G.B. Pant University of Agri. & Tech., Pantnagar, India

Prof. Jitendra Kwatra, Ph.D., Director, Extension Education, G.B. Pant University of Agri. & Tech., Pantnagar, India

Prof. S.S. Gupta, Ph.D., Dean, College of Technology, G.B. Pant University of Agri. & Tech., Pantnagar, India

Prof. D. Kumar, Ph.D., Dean, College of Veterinary & Animal Sciences, G.B. Pant University of Agri. & Tech., Pantnagar, India

Prof. Alka Goel, Ph.D., Dean, College of Community Science, G.B. Pant University of Agri. & Tech., Pantnagar, India

Prof. R.S. Jadoun, Ph.D., Dean, College of Agribusiness Management, G.B. Pant University of Agri. & Tech., Pantnagar, India

Prof. Lokesh Varshney, Ph.D., Dean, College of Post Graduate Studies, G.B. Pant University of Agri. & Tech., Pantnagar, India

Prof. Avdhesh Kumar, Ph.D., Dean, College of Fisheries, G.B. Pant University of Agri. & Tech., Pantnagar, India

Prof. Subhash Chandra, Ph.D., Dean, College of Agriculture, G.B. Pant University of Agri. & Tech., Pantnagar, India

Prof. Ramesh Chandra Srivastava, Ph.D., Dean, College of Basic Sciences & Humanities, G.B.P.U.A.T., Pantnagar, India

EDITORIAL BOARD

Members

A.K. Misra, Ph.D., Ex-Chairman, Agricultural Scientists Recruitment Board, Krishi Anusandhan Bhavan I, New Delhi, India & Ex-Vice Chancellor, G.B. Pant University of Agriculture & Technology, Pantnagar

Anand Shukla, Director, Reefberry Foodex Pvt. Ltd., Veraval, Gujarat, India

Anil Kumar, Ph.D., Director, Education, Rani Lakshmi Bai Central Agricultural University, Jhansi, India

Ashok K. Mishra, Ph.D., Kemper and Ethel Marley Foundation Chair, W.P. Carey Business School, Arizona State University, U.S.A.

Binod Kumar Kanaujia, Ph.D., Professor, School of Computational and Integrative Sciences, Jawahar Lal Nehru University, New Delhi, India

D. Ratna Kumari, Ph.D., Associate Dean, College of Community / Home Science, PJTSAU, Hyderabad, India

Deepak Pant, Ph.D., Separation and Conversion Technology, Flemish Institute for Technological Research (VITO), Belgium

Desirazu N. Rao, Ph.D., Honorary Professor, Department of Biochemistry, Indian Institute of Science, Bangalore, India

G.K. Garg, Ph.D., Ex-Dean, College of Basic Sciences & Humanities, G.B. Pant University of Agri. & Tech., Pantnagar, India

Humnath Bhandari, Ph.D., IRRI Representative for Bangladesh, Agricultural Economist, Agrifood Policy Platform, Philippines

Indu S. Sawant, Ph.D., Principal Scientist, ICAR National Research Centre for Grapes, Pune, India

Kuldeep Singh, Ph.D., Director, ICAR - National Bureau of Plant Genetic Resources, New Delhi, India

Muneshwar Singh, Ph.D., Ex-Project Coordinator AICRP- LTFE, ICAR, Indian Institute of Soil Science, Bhopal, India

Omkar, Ph.D., Professor (Retd.), Department of Zoology, University of Lucknow, India

P.C. Srivastav, Ph.D., Professor (Retd.), Department of Soil Science, G.B. Pant University of Agriculture and Technology, Pantnagar, India

Prashant Srivastava, Ph.D., Soil Contaminant Chemist, CSIRO, Australia

Puneet Srivastava, Ph.D., Director, Water Resources Center, Butler-Cunningham Eminent Scholar, Professor, Biosystems Engineering, Auburn University, United States

R.K. Singh, Ph.D., Ex-Director & Vice Chancellor, ICAR-Indian Veterinary Research Institute, Izatnagar, U.P., India

Ramesh Kanwar, Ph.D., Charles F. Curtiss Distinguished Professor of Water Resources Engineering, Iowa State University, U.S.A.

S.N. Maurya, Ph.D., Professor (Retired), Department of Gynaecology & Obstetrics, G.B. Pant University of Agri. & Tech., Pantnagar, India

Sham S. Goyal, Ph.D., Professor Emeritus, Faculty of Agriculture and Environmental Sciences, University of California, Davis, U.S.A.

Umesh Varshney, Ph.D., Honorary Professor, Department of Microbiology and Cell Biology, Indian Institute of Science, Bangalore, India

V.D. Sharma, Ph.D., Dean Life Sciences, SAI Group of Institutions, Dehradun, India

V.K. Singh, Ph.D., Director, ICAR-Central Research Institute for Dryland Agriculture, Hyderabad, India

Vijay P. Singh, Ph.D., Distinguished Professor, Caroline and William N. Lehrer Distinguished Chair in Water Engineering, Department of Biological and Agricultural Engineering, Texas A & M University, U.S.A.

Nikitin Georgy, Ph.D., Vice-Rector cum Professor St. Petersburg State University of Veterinary Medicine St. Petersburg, Russia

Sanjay Jambhulkar, Ph.D., Ex-Scientific Officer 'H' Head Mutation Breeding Section Bhabha Atomic Research Centre, Mumbai Professor, Homi Bhabha National Institute, Mumbai

Vijay Kubsad, Ph.D., Adjunct Professor Washington State University USA

Kiran Tota-Maharaj, Ph.D., Professor Royal Agricultural University (RAU) Cirencester, Gloucestershire GL76JS England, UK

Ravindra P. Veeranna, Ph.D., Dean, Xavier University, College of Veterinary Sciences, Aruba

Javed Akhtar, Ph.D., University of Technology and Applied Sciences, Muscat

Basant Maheshwari, Ph.D., Western Sydney University, Australia

Vijay P. Singh, Ph.D. Professor Emeritus, Texas A&M University College station, Texas, USA

Sanjay Sharma, Ph.D. Director, Uttarakhand Council of Biotechnology

Krishna Nemali, Ph.D. Controlled Environment Agriculture Extension Specialist

Editor-in-Chief

K.P. Raverkar, Professor, G.B. Pant University of Agriculture and Technology, Pantnagar, India

Managing Editor

S.K. Guru, Ph.D., Professor, G.B. Pant University of Agriculture and Technology, Pantnagar, India

Assistant Managing Editor

Jyotsna Yadav, Ph.D., Research Editor, Directorate of Research, G.B. Pant University of Agriculture and Technology, Pantnagar, India

Technical Manager

S.D. Samantaray, Ph.D., Professor & Head, Department of Computer Engineering, G.B. Pant University of Agriculture and Technology, Pantnagar, India

Development

Dr. S.D. Samantaray, Professor & Head

Brijesh Dumka, Developer & Programmer

CONTENTS

Indian mustard: Phytoremediation tool for soil health management (Review)	1
M.L. DOTANIYA, C.K. DOTANIYA, KULDEEP KUMAR, JAKIR HUSSAIN, R.K. DOUTANIYA, H.M. MEENA and MANJU LATA	
From profile to problems: A study on socio-economic status and constraints of rural women in dairy farming	14
ADITI PATHAK and ARPITA SHARMA KANDPAL	
Diversification and intensification in rice–wheat cropping system for enhancing productivity under <i>Tarai</i> zone of Uttarakhand	20
PESALA SANTHA KUMAR, DINESH KUMAR SINGH, SUBHASH CHANDRA, ROHITASH SINGH and AJEET PRATAP SINGH	
Sensor based nitrogen application impact on productivity, economics and nutrient uptake in maize	25
AMIT BHATNAGAR and AJAY KUMAR	
Association of germination and seedling vigour attributes with green forage yield in oat (<i>Avena sativa</i> L.) genotypes through correlation and path coefficient analysis	31
DEVYANI RAMOLA, AKRITI BALLABH SANSKRITI JOSHI, MANISH MEHRA and BIRENDRA PRASAD	
Genetic divergence and yield trait impact in French bean (<i>Phaseolus vulgaris</i> L.)	43
BHAWANA MAMGAIN, B.P. NAUTIYAL, S.C. PANT, VAISHALI THAPA, HEIGRUJAM RIYASANA and MAYANK	
Arabic-gum based edible coatings preserving the post-harvest quality of Guava fruits	48
ESHA TEWARI and OMVEER SINGH	
Efficacy of catechol and plant growth regulators on the vegetative growth and flowering of African Marigold (<i>Tagetes erecta</i> L. cv. Pusa Narangi Gaiinda)	59
SURYANSH SRIVASTAVA, AFSAR AHMAD, MD. RAMJAN and KHAN BILAL MUKHTAR AHMED	
Host response of diverse pulse genotypes to root-knot nematode (<i>Meloidogyne incognita</i>)	67
SAROJ YADAV, EKTA RATHORE and ANIL KUMAR	
Effect of egg size on egg quality traits, fertility, hatchability and day-old chick weight	75
ISLAM UDDIN SHEIKH, AZMAT ALAM KHAN, ARNAB JYOTI KALITA, SHEIKH ADIL HAMID, IRFAN AKRAM BABA, HENNA HAMADANI, BUSHRA ZAFFER and INSHA AFZAL	
Pathomorphological studies on verminous pneumonia in a Gaddi goat	81
SWASTI SHARMA, SONALI MISHRA and R.D. PATIL	
Pathological findings in congestive heart failure in dog: A case report	85
MUNISH BATRA and STUTI SINGH	

Histopathological and haematological evaluation of the ameliorative effect of piperine on quinalphos-induced subacute toxicity in Swiss Albino Mice PREETI SINGH, VINOD KUMAR, PREETI BAGRI and DEEPIKA LATHER	89
Phytochemical screening and antioxidant activities of locally available medicinal plants PREETI BAGRI, VIJEYTA TIWARI, ARCHANA LOHIYA and VINOD KUMAR	99
Natural Farming Inputs (NFIs) as sustainable alternatives for enhancing the growth and development of Chickpea (<i>Cicer arietinum</i> L.) SHAIFALI and A.K. SHARMA	108
Integrating Bayesian Calibration, Hierarchical Spatial Regression, and Structural–Temporal Models to Explain and Predict Wheat Yield in the Tarai–Bhabar Belt of Uttarakhand, India NEHA BHATT and VINOD KUMAR	114
Phytochemical characterization and immuno-antioxidant potential of <i>Moringa oleifera</i> leaf extract in chicken lymphocytes ANAMITRA RUJ, SONU AMBWANI and LATA PALIWAL	128
Biochemical changes associated with triclosan exposure in the freshwater fish <i>Cyprinus carpio</i> MADHU SHARMA, YASHPAL, RIJUL RANA, DEEKSHA, DEEPIKA THAKUR and TARANG KUMAR SHAH	140
Assessment of awareness among respondents towards Pradhan Mantri Jan Dhan Yojana: A study in Haryana MEGHA GOYAL, SUMAN GHALAWAT, MINAKSHI, ATUL DHINGRA and MINI	148
Youth perception and constraints in agri-entrepreneurship: A factor analytic study JYOTI, SUMAN GHALAWAT, MEGHA GOYAL, NEELAM KAUSHAL, SUNIL KUMAR and VIKRANT HOODA	156
Design of Object Detection Model Using Deep Learning for Autonomous Driving in Adverse Weather Conditions: A GAN-Based Restoration and Depth-Aware Fusion Framework MUMUKSHU BHATT, SUBODH PRASAD, RAJESH SINGH, ASHOK KUMAR and POOJA TAMTA	165
A novel mechanical purification approach for <i>Grewia optiva</i> Fibre-Reinforced epoxy composites: Effect of fibre geometry on structural performance DEEPA SINGH, NEERAJ BISHT, TARUN KUMAR SAXENA and KAMAL RAWAT	172
Evaluating Long-Term rainfall and temperature trends across the Shipra River basin POONAM SINGH, MOHAN LAL, DHEERAJ KUMAR, P. K. SINGH and GOVIND VERMA	184
Rainfall-runoff modelling using soft computing techniques for various watersheds of Madhya Pradesh, India SEEMA KUSHWAHA and PRAVENDRA KUMAR	196
Prevalence and associated risk factors of functional constipation among the selected adult population of Pantnagar, U.S. Nagar, Uttarakhand, India JYOTI SINGH, NEETU DOBHAI, ARCHANA KUSHWAHA, N.C SHAHI and SABBU SANGEETA	204
Tracking nutritional transitions: A comparative study of child malnutrition (0–5 years) trends across districts of Madhya Pradesh SHALINI CHAKRABORTY	212
Determinants influencing tourist selection of rural tourism destinations using Garrett Ranking Technique: Evidence from Supi village, Uttarakhand NEHA PANDEY and KAMAL UPRETI	221

Diversification and intensification in rice–wheat cropping system for enhancing productivity under *Tarai* zone of Uttarakhand

PESALA SANTHA KUMAR*, DINESH KUMAR SINGH, SUBHASH CHANDRA, ROHITASH SINGH and AJEET PRATAP SINGH¹

Department of Agronomy, ¹Department of Soil Science, College of Agriculture, G. B. Pant University of Agriculture and Technology, Pantnagar- 263145 (U. S. Nagar, Uttarakhand)

**Corresponding author's email id: shanthpesala1834@gmail.com*

ABSTRACT: The field experiment was conducted during 2021-2022 at Norman E. Borlaug Crop Research Centre, G.B Pant University of Agriculture and Technology, Pantnagar, U.S Nagar, Uttarakhand, India to evaluate the impact of diversification and intensification in rice-wheat cropping system (RWCS) on productivity and profitability. The experimental field soil was loam in texture, high in organic carbon, low in available nitrogen, high in phosphorus and medium in potassium with neutral pH. Ten cropping sequences were evaluated in a randomized block design with three replications. The study formed part of a long-term trial initiated in 2017–18 under AICRP–IFS. Results revealed that the rice–vegetable pea–mentha cropping sequence recorded the highest system productivity (239.7 q REY/ha), production efficiency (66.4 kg REY/ha/day) and relative production efficiency (148.1%), outperforming the conventional rice–wheat–fallow sequence. Napier grass + fodder cowpea–napier grass + berseem–napier grass + fodder cowpea cropping recorded the highest land use efficiency (99.4%), where as nutrient productivity was recorded highest in rice–vegetable pea–black gram and rice–vegetable pea–mentha cropping systems mainly due to higher productivity of vegetable pea (133.4 and 121.5 q REY/ha, respectively). The presence of high-value crops (mentha and vegetable pea) contributed to the superior performance of diversified systems. Overall, diversification of conventional RWCS with legumes, vegetables and fodder crops substantially enhanced productivity, system efficiency, resource-use efficiency and economic profitability. Under the *Tarai* conditions of Uttarakhand, the rice–vegetable pea–mentha sequence proved to be the most productive and remunerative option among the tested cropping systems.

Keywords: Equivalent yield, Land use efficiency, Nutrient productivity, Production efficiency, System productivity

The rice–wheat cropping system (RWCS) is one of the world's largest agricultural production systems, covering nearly 26 million ha across the Indo-Gangetic Plains (IGP) of South Asia (Shweta and Malik, 2017). In India, RWCS occupies about 40% of the IGP's cultivated area and plays a major role in national food security. However, the system faces several sustainability challenges, including declining groundwater levels, nutrient imbalance, soil degradation, salinity, pest and disease resurgence, environmental pollution and reduced farm profitability (Parihar *et al.*, 2016).

Crop diversification has emerged as an effective strategy to address these challenges by improving food and nutritional security, enhancing income, promoting efficient use of land and water resources. Integrating legumes into the rice–wheat system

enhances productivity per unit resources, conserves groundwater and soil moisture and reduces input costs through biological nitrogen fixation. Diversification with maize, pulses, oilseeds and high-value cash crops such as potato, mentha and broccoli helps to reduce climatic and market risks while improving system productivity and profitability.

Inclusion of fodder crops such as sorghum and Napier grass can further strengthen integrated farming systems by providing nutrient-rich green fodder with high dry matter yield. Overall, diversification offers a viable approach to overcome the limitations of cereal–cereal rotations and enhance the productivity, profitability and sustainability of RWCS, particularly under irrigated conditions such as the *Tarai* region.

MATERIALS AND METHODS

A long-term experiment on diversification and intensification of cropping systems was conducted from 2017–18 to 2021–22 under the All India Coordinated Research Programme on Integrated Farming Systems (AICRP–IFS). The present study reports results from the 2021–22 cropping cycle, carried out at E-2 block of the Norman E. Borlaug Crop Research Centre, Govind Ballabh Pant University of Agriculture and Technology, Pantnagar, Udham Singh Nagar, Uttarakhand, India. The site is located at 29° N latitude and 79.3° E longitude, at an altitude of 243.83 m above mean sea level, within the *Tarai* belt of the Shivalik foothills. This region has a subtropical climate characterized by hot summers (up to 40°C) and cold winters (occasionally below 0°C), with an average annual rainfall of about 1450 mm, of which 80–90% is received during June to September. The soils are calcareous, medium to moderately coarse-textured and classified under the Mollisol order.

The experiment was laid out to evaluate the effects of diversification and intensification within the rice–wheat cropping system. During the rainy (*kharif*) season, crops such as rice, black gram, maize and sorghum were sown in the first week of July according to monsoon onset. Field preparation for both *kharif* and *rabi* crops consisted of one deep ploughing with a mouldboard plough followed by two cross harrowings and planking. Post-rainy (*rabi*) crops were sown during the first fortnight of November.

Details of the experimental treatments and the cultural practices adopted during the study are presented in Table 1 and Table 2. Standard agronomic practices were followed for each crop as per recommended packages.

Rice equivalent yield (REY), production efficiency, relative production efficiency, land-use efficiency and nutrient productivity were computed using standard formulae. Data were analyzed using ANOVA for randomized block design. Treatment means were compared using LSD at 5% significance.

1. Rice Equivalent Yield (REY)

$$REY = \frac{Y_i \times P_i}{P_r}$$

Where,

Y_i = Yield of *i*th crop (kg ha⁻¹);

P_i = Price of *i*th crop (Rs kg⁻¹);

P_r = Price of rice (Rs kg⁻¹)

2. Production Efficiency (PE)

$$PE = \frac{REY}{D}$$

Where,

REY_t = Rice equivalent yield (kg ha⁻¹);

D = Duration of cropping system (days)

3. Relative Production Efficiency (RPE)

$$RPE(\%) = \frac{REY_t - REY_c}{REY_c} \times 100$$

Where,

REY_t = REY of treatment;

REY_c = REY of control (rice–wheat)

4. Land Use Efficiency (LUE)

$$LUE(\%) = \frac{DO}{365} \times 100$$

Where, DO = Days land occupied

5. Nutrient Productivity (NP)

$$NP = \frac{REY}{N + P + K}$$

Where, N, P, K = Total nutrients applied (kg ha⁻¹)

RESULTS AND DISCUSSION

System Productivity

Significant variation in system productivity was observed among the treatments. The rice–vegetable pea–mentha cropping sequence recorded the highest

productivity (239.7 q REY/ha), followed by rice–vegetable pea–black gram (223.7 q REY/ha) and basmati rice–potato–maize (cob) (218.8 q REY/ha). The above cropping systems produced 2.47, 2.31 and 2.26 times higher system productivity respectively than the traditional rice–wheat cropping system (96.7 q REY/ha). The lowest productivity (29.0 q REY/ha) was noted in the fodder-based napier grass + fodder cowpea – napier grass + berseem – napier grass + fodder cowpea cropping system.

Higher productivity in rice–vegetable pea–mentha, rice–vegetable pea–black gram and basmati rice–potato–maize (cob) resulted primarily from the inclusion of short-duration and high-yielding crops such as vegetable in rice–vegetable pea–mentha (121.5 q REY/ha) and rice–vegetable pea–black gram (133.4 q REY/ha) and potato in basmati rice–potato–maize (cob) cropping system (89.4 q REY/ha). These findings corroborate the results of Singh *et al.* (2013), who reported that inclusion of vegetables enhances system productivity due to their higher production potential. The lower values observed in fodder-based systems across productivity parameters are primarily due to the lower economic conversion efficiency of fodder crops, which produce high biomass but comparatively low marketable yield and price, thereby reducing rice equivalent yield and overall system productivity, as also reported by Li *et al.* (2024).

Production Efficiency

Production efficiency followed a similar pattern,

rice–vegetable pea–mentha cropping sequence recorded the highest value (66.4 kg REY/ha/day), followed by rice–vegetable pea–black gram (61.8 kg REY/ha/day) and basmati rice–potato–maize (cob) (59.8 kg REY/ha/day). These systems achieved 2.51, 2.34 and 2.26 times greater production efficiency than the rice–wheat system (26.4 kg REY/ha/day). The lowest value (7.9 kg REY/ha/day) occurred in napier grass + fodder cowpea – napier grass + berseem – napier grass + fodder cowpea.

The superior performance of rice–vegetable pea–mentha, rice–vegetable pea–black gram and basmati rice–potato–maize (cob) can be attributed to higher daily productivity from vegetable pea and potato, which yield substantially within a shorter duration, thereby increasing output per unit time. Similar conclusions were drawn by Prasad *et al.* (2013), who noted that vegetable-based sequences showed higher production efficiency due to their high yield potential within a short growth period.

Relative Production Efficiency

Relative production efficiency (RPE) was highest (148.1%) in rice–vegetable pea–mentha, followed by rice–vegetable pea–black gram (131.6%). The lowest values (–70%) occurred in fodder-based system such as napier grass + fodder cowpea – napier grass + berseem – napier grass + fodder cowpea and black gram – wheat – vegetable cowpea (–11.1%). The higher RPE in rice–vegetable pea–mentha and rice–vegetable pea–black gram resulted from the inclusion of vegetables and legumes during the zaid season, leading to substantially higher system productivity compared with the fallow period of the

Table 1: Treatment details of experiment

Treatments	Kharif	Rabi	Zaid
T ₁	Rice	Wheat	Fallow
T ₂	Rice	Vegetable pea	Mentha
T ₃	Blackgram	Wheat	Vegetable cowpea
T ₄	Rice	Vegetable pea	Blackgram
T ₅	Maize	Yellow sarson	Blackgram
T ₆	Rice	Yellow sarson	Grain Cowpea
T ₇	Multicut sorghum	Berseem+ oat	Maize+ Cowpea
T ₈	Napiergrass+ Fodder cowpea	Napiergrass+ Berseem	Napiergrass+ Fodder cowpea
T ₉	Basmati Rice	Potato	Maize (Cobs)
T ₁₀	Maize (Cobs)	Broccoli	Okra

Table 2: Agronomic practices followed in different crops

Crop	Seedrate (kg/ha)	Fertilizer rate (NPK kg/ha)	Spacing (cmxcm)	Cultivar/ Variety
Rice	35 (50)	150:60:40	20 x10	HKR-47
Wheat	100 (125)	150:60:40	22.5	UP-2784
Maize	10	120:60:40	60 x15	Suvarna
Mentha	400 kg suckers	120:60:40	60	Him Kranti
Potato	2000	200:100:100	60 x15	Kufri Himsona
Veg. pea	100	20:60:40	30 x5	Arkel
Black Gram	20	20:40:20	30 x10	Pant Urd-31
Yellow sarson	5	60:30:30	30 x 15	PPS-1
Cowpea	10	20:40:0	35 x5	Kashi Kanchan
Broccoli	28000 seedlings	120:60:40	60 x60	Perasio (Hybrid)
Okra	12	120:60:40	60 x15	Kashi Pragati
Sorghum	25	120:60:40	Broadcasting	PC-5
Berseem	25	20:60:40	30 x5	Mescavi
Oat	100	120:60:40	23	UPO-212
Basmati Rice	15	150:60:40	20 x10	PUSA-1509

Table 3: Effect of diversification and intensification of cropping system on system productivity, production efficiency relative production efficiency, land use efficiency (LUE) and nutrient productivity

Treatments	System Productivity (qREY/ha)	Production efficiency (kgREY/ha/day)	Relative production efficiency(%)	Crop growing periods (days)	LUE (%)	Nutrient productivity (kgREY/ kgNPK)
T1	96.7	26.4	--	259	70.9	19.3
T2	239.7	66.4	148.1	315	86.3	40.6
T3	85.9	23.5	-11.1	325	89.0	20.9
T4	223.7	61.8	131.6	293	80.2	49.7
T5	114.4	31.5	18.4	324	88.7	27.2
T6	121.3	33.2	25.5	341	93.4	28.2
T7	154.3	42.5	59.7	353	96.7	29.1
T8	29.0	7.90	-70.0	363	99.4	5.7
T9	218.8	59.8	126.4	305	83.5	24.3
T10	170.2	46.4	76.1	291	79.7	25.8
SEm±	2.06	0.35	27.8	-	-	0.39
CD (p=0.05)	6.12	1.03	82.8	-	-	1.16

traditional rice–wheat system. These observations align with Mishra *et al.* (2007), who reported that addition of short-duration legumes, vegetables and medicinal crops during the summer season improves overall system productivity.

Land-Use Efficiency

Among the different systems, napier grass + fodder cowpea – napier grass + berseem – napier grass + fodder cowpea recorded the highest (99.4%) land-use efficiency followed by multicut sorghum – berseem – oat – maize + fodder cowpea (96.7%). In contrast, the rice–wheat–fallow system exhibited the lowest land-use efficiency (70.9%). Higher LUE in fodder-based systems was mainly due to continuous land occupancy (363 days in napier grass + fodder

cowpea – napier grass + berseem – napier grass + fodder cowpea and 353 days in multicut sorghum – berseem – oat – maize + fodder cowpea). The higher land-use efficiency observed in fodder-based systems was primarily due to continuous land occupancy throughout the year, as these systems involve perennial and multi-cut fodder crops with negligible fallow period, resulting in maximum temporal utilization of land resources, in line with the findings of Li *et al.* (2024).

Nutrient Productivity

Similarly, variation was also observed in nutrient productivity across systems. Nutrient productivity was highest (49.7 kg REY/ kg NPK) in rice–vegetable pea–black gram cropping sequence

followed by rice–vegetable pea–mentha, mainly due to higher productivity of vegetable pea (133.4 and 121.5 q REY/ha, respectively) at a relatively low nutrient application rate (20:40:0 NPK). Legumes, being nutrient-efficient and capable of biological nitrogen fixation, provided higher yield per unit nutrient applied compared with nutrient-exhaustive cereals. Similar findings were reported by Kumar *et al.* (2023), who observed higher nutrient productivity in legume-based cropping sequences.

CONCLUSION

The study demonstrates that diversification and intensification of the rice–wheat system significantly improves productivity and economic returns. Among the evaluated systems, the rice–vegetable pea–mentha cropping sequence emerged as the most efficient and profitable. Therefore, inclusion of high-value vegetable crops in the rice-based system is recommended as a viable strategy to enhance farm profitability and resource-use efficiency under *Tarai* conditions.

REFERENCES

- Kumar, S., Singh, R. and Patel, A. (2023). Soil health improvement through crop diversification in rice-based systems. *J. Soil Sci. Plant Nutr.*, 23: 1123–1135.
- Li, H., Zhang, Q. and Chen, J. (2024). Impact of legumes on soil fertility and nutrient cycling. *Agron. J.*, 116(3): 1456–1468.
- Mishra, M. M., Nanda, S. S., Mohanty, M., Pradhan, K. C. and Mishra, S. S. (2007). Crop diversification under rice-based cropping system in western Orissa. In: *Extended summaries 3 National Symposium on Integrated Farming Systems* at Agricultural Research Station, Durgapura, Jaipur, during October 26-28. 258p.
- Parihar, C.M., Yadav, M.R., Jat, S.L., Singh, A.K., Kumar, B., Pradhan, S. and Chakraborty, D. (2016). Long-term effect of conservation agriculture in maize rotations on total organic carbon, physical and biological properties of a sandy loam soil in north-western Indo-Gangetic Plains. *Soil Till Res.*, 161:116-28.
- Prasad, D., Yadava, M. S. and Singh, C. S. (2013). Diversification of rice (*Oryza sativa*)-based cropping systems for higher productivity, profitability and resource-use efficiency under irrigated ecosystem of Jharkhand. *Indian J. Agron.*, 58(3): 264-270.
- Shweta, M. M. and Malik, M. (2017). Improving wheat productivity in rice-wheat cropping system through crop establishment methods. *Int. J. Pure. Appl. Biosci.*, 5(3): 575-578.
- Singh, R. K., Bohra, J. S., Nath, T. and Srivastava, R. K. (2013). Effect of varying level of diversification and intensification of rice-wheat cropping system on the production potential and nutrient balance of soil in Indo-Gangetic Plain of India. *Agric.*, 10(2): 9-16.

Received: April 17, 2026

Accepted: April 28, 2026