Print ISSN : 0972-8813
e-ISSN : 2582-2780

[Vol. 24(1) January-April 2026]

Pantnagar Journal of Research

(Formerly International Journal of Basic and Applied Agricultural Research ISSN : 2349-8765)

G.B. Pant University of Agriculture & Technology
Pantnagar, U.S. Nagar; Uttarakhand, Website : gbpuat.res.in/PJR

ADVISORY BOARD

Patron

Prof. Shivendra Kumar Kashyap, Ph.D., Vice-Chancellor, G.B. Pant University of Agriculture and Technology, Pantnagar, India

Members

Prof. S. K. Verma, Ph.D., Director Research, G.B. Pant University of Agri. & Tech., Pantnagar, India
Prof. Jitendra Kwatra, Ph.D., Director, Extension Education, G.B. Pant University of Agri. & Tech., Pantnagar, India
Prof. S.S. Gupta, Ph.D., Dean, College of Technology, G.B. Pant University of Agri. & Tech., Pantnagar, India
Prof. D. Kumar, Ph.D., Dean, College of Veterinary & Animal Sciences, G.B. Pant University of Agri. & Tech., Pantnagar, India
Prof. Alka Goel, Ph.D., Dean, College of Community Science, G.B. Pant University of Agri. & Tech., Pantnagar, India
Prof. R.S. Jadoun, Ph.D., Dean, College of Agribusiness Management, G.B. Pant University of Agri. & Tech., Pantnagar, India
Prof. Lokesh Varshney, Ph.D., Dean, College of Post Graduate Studies, G.B. Pant University of Agri. & Tech., Pantnagar, India
Prof. Avdhesh Kumar, Ph.D., Dean, College of Fisheries, G.B. Pant University of Agri. & Tech., Pantnagar, India
Prof. Subhash Chandra, Ph.D., Dean, College of Agriculture, G.B. Pant University of Agri. & Tech., Pantnagar, India
Prof. Ramesh Chandra Srivastava, Ph.D., Dean, College of Basic Sciences & Humanities, G.B.P.U.A.T., Pantnagar, India

EDITORIAL BOARD

Members

A.K. Misra, Ph.D., Ex-Chairman, Agricultural Scientists Recruitment Board, Krishi Anusandhan Bhavan I, New Delhi, India & Ex-Vice Chancellor, G.B.Pant University of Agriculture & Technology, Pantnagar
Anand Shukla, Director, Reefberry Foodex Pvt. Ltd., Veraval, Gujarat, India
Anil Kumar, Ph.D., Director, Education, Rani Lakshmi Bai Central Agricultural University, Jhansi, India
Ashok K. Mishra, Ph.D, Kemper and Ethel Marley Foundation Chair, W P Carey Business School, Arizona State University, U.S.A
Binod Kumar Kanaujia, Ph.D., Professor, School of Computational and Integrative Sciences, Jawahar Lal Nehru University, New Delhi, India
D. Ratna Kumari, Ph.D., Associate Dean, College of Community / Home Science, PJTSAU, Hyderabad, India
Deepak Pant, Ph.D., Separation and Conversion Technology, Flemish Institute for Technological Research (VITO), Belgium
Desirazu N. Rao, Ph.D., Honorary Professor, Department of Biochemistry, Indian Institute of Science, Bangalore, India
G.K. Garg, Ph.D., Ex-Dean, College of Basic Sciences & Humanities, G.B.Pant University of Agric. & Tech., Pantnagar, India
Humnath Bhandari, Ph.D., IRRI Representative for Bangladesh, Agricultural Economist, Agrifood Policy Platform, Philippines
Indu S. Sawant, Ph.D., Principal Scientist, ICAR National Research Centre for Grapes, Pune, India
Kuldeep Singh, Ph.D. , Director, ICAR - National Bureau of Plant Genetic Resources, New Delhi, India
Muneshwar Singh, Ph.D., Ex-Project Coordinator AICRP- LTFE, ICAR, Indian Institute of Soil Science, Bhopal, India
Omkar, Ph.D., Professor (Retd.), Department of Zoology, University of Lucknow, India
P.C. Srivastav, Ph.D., Professor (Retd.), Department of Soil Science, G.B. Pant University of Agriculture and Technology, Pantnagar, India
Prashant Srivastava, Ph.D., Soil Contaminant Chemist, CSIRO, Australia
Puneet Srivastava, Ph.D., Director, Water Resources Center, Butler-Cunningham Eminent Scholar, Professor, Biosystems Engineering, Auburn University, United States
R.K. Singh, Ph.D., Ex-Director & Vice Chancellor, ICAR-Indian Veterinary Research Institute, Izatnagar, U.P., India
Ramesh Kanwar, Ph.D., Charles F. Curtiss Distinguished Professor of Water Resources Engineering, Iowa State University, U.S.A.
S.N. Maurya, Ph.D., Professor (Retired), Department of Gynaecology & Obstetrics, G.B. Pant University of Agric. & Tech., Pantnagar, India
Sham S. Goyal, Ph.D., Professor Emeritus, Faculty of Agriculture and Environmental Sciences, University of California, Davis, U.S.A.
Umesh Varshney, Ph.D., Honorary Professor, Department of Microbiology and Cell Biology, Indian Institute of Science, Bangalore, India
V.D. Sharma, Ph.D., Dean Life Sciences, SAI Group of Institutions, Dehradun, India
V.K. Singh, Ph.D., Director, ICAR-Central Research Institute for Dryland Agriculture, Hyderabad, India
Vijay P. Singh, Ph.D., Distinguished Professor, Caroline and William N. Lehrer Distinguished Chair in Water Engineering, Department of Biological and Agricultural Engineering, Texas A & M University, U.S.A.
Nikitin Georgy, Ph.D., Vice-Rector cum Professor St. Petersberg State University of Veterinary Medicine St. Petersberg, Russia
Sanjay Jambhulkar, Ph.D., Ex-Scientific Officer 'H' Head Mutation Breeding Section Bhabha Atomic Research Centre, Mumbai Professor, Homi Bhabha National Institute, Mumbai
Vijay Kubsad, Ph.D., Adjunct Professor Washington State University USA
Kiran Tota-Maharaj, Ph.D., Professor Royal Agricultural University (RAU) Cirencester, Gloucestershire GL76JS England, UK
Ravindra P. Veeranna, Ph.D., Dean, Xavier University, College of Veterinary Sciences, Aruba
Javed Akhtar, Ph.D., University of Technology and Applied Sciences, Muscat
Basant Maheshwari, Ph.D., Western Sydney University, Australia
Vijay P. Singh, Ph.D. Professor Emeritus, Texas A&M University College station, Texas, USA
Sanjay Sharma, Ph.D. Director, Uttarakhand Council of Biotechnology
Krishna Nemali, Ph.D. Controlled Environment Agriculture Extension Specialist

Editor-in-Chief

K.P. Raverkar, Professor, G.B. Pant University of Agriculture and Technology, Pantnagar, India

Managing Editor

S.K. Guru, Ph.D., Professor, G.B. Pant University of Agriculture and Technology, Pantnagar, India

Assistant Managing Editor

Jyotsna Yadav, Ph.D., Research Editor, Directorate of Research, G.B. Pant University of Agriculture and Technology, Pantnagar, India

Technical Manager

S.D. Samantaray, Ph.D., Professor & Head, Department of Computer Engineering, G.B. Pant University of Agriculture and Technology, Pantnagar, India

Development

Dr. S.D. Samantaray, Professor & Head
Brijesh Dumka, Developer & Programmer

PANTNAGAR JOURNAL OF RESEARCH

Vol. 24(1) January-April, 2026

CONTENTS

Indian mustard: Phytoremediation tool for soil health management (Review) **1**
M.L. DOTANIYA, C.K. DOTANIYA, KULDEEP KUMAR, JAKIR HUSSAIN, R.K. DOUTANIYA, H.M. MEENA and MANJU LATA

From profile to problems: A study on socio-economic status and constraints of rural women in dairy farming **14**
ADITI PATHAK and ARPITA SHARMA KANDPAL

Diversification and intensification in rice–wheat cropping system for enhancing productivity under *Tarai* zone of Uttarakhand **20**
PESALA SANTHA KUMAR, DINESH KUMAR SINGH, SUBHASH CHANDRA, ROHITASH SINGH and AJEET PRATAP SINGH

Sensor based nitrogen application impact on productivity, economics and nutrient uptake in maize **25**
AMIT BHATNAGAR and AJAY KUMAR

Association of germination and seedling vigour attributes with green forage yield in oat (*Avena sativa* L.) genotypes through correlation and path coefficient analysis **31**
DEVYANI RAMOLA, AKRITI BALLABH SANSKRITI JOSHI, MANISH MEHRA and BIRENDRA PRASAD

Genetic divergence and yield trait impact in French bean (*Phaseolus vulgaris* L.) **43**
BHAWANA MAMGAIN, B.P. NAUTIYAL , S.C. PANT, VAISHALI THAPA, HEIGRUJAM RIYASANA and MAYANK

Arabic-gum based edible coatings preserving the post-harvest quality of Guava fruits **48**
ESHA TEWARI and OMVEER SINGH

Efficacy of catechol and plant growth regulators on the vegetative growth and flowering of African Marigold (*Tagetes erecta* L. cv. Pusa Narangi Gainda) **59**
SURYANSH SRIVASTAVA, AFSAR AHMAD, MD. RAMJAN and KHAN BILAL MUKHTAR AHMED

Host response of diverse pulse genotypes to root-knot nematode (*Meloidogyne incognita*) **67**
SAROJ YADAV, EKTA RATHORE and ANIL KUMAR

Effect of egg size on egg quality traits, fertility, hatchability and day-old chick weight **75**
ISLAM UDDIN SHEIKH, AZMAT ALAM KHAN, ARNAB JYOTI KALITA, SHEIKH ADIL HAMID, IRFAN AKRAM BABA, HENNA HAMADANI, BUSHRA ZAFFER and INSHA AFZAL

Pathomorphological studies on verminous pneumonia in a Gaddi goat **81**
SWASTI SHARMA, SONALI MISHRA and R.D. PATIL

Pathological findings in congestive heart failure in dog: A case report **85**
MUNISH BATRA and STUTI SINGH

Histopathological and haematological evaluation of the ameliorative effect of piperine on quinalphos-induced subacute toxicity in Swiss Albino Mice 89
PREETI SINGH, VINOD KUMAR, PREETI BAGRI and DEEPIKA LATHER

Phytochemical screening and antioxidant activities of locally available medicinal plants 99
PREETI BAGRI, VIJEYTA TIWARI, ARCHANA LOHIYA and VINOD KUMAR

Natural Farming Inputs (NFIs) as sustainable alternatives for enhancing the growth and development of Chickpea (*Cicer arietinum* L.) 108
SHAIFALI and A.K. SHARMA

Integrating Bayesian Calibration, Hierarchical Spatial Regression, and Structural–Temporal Models to Explain and Predict Wheat Yield in the Tarai–Bhabar Belt of Uttarakhand, India 114
NEHA BHATT and VINOD KUMAR

Phytochemical characterization and immuno-antioxidant potential of *Moringa oleifera* leaf extract in chicken lymphocytes 128
ANAMITRA RUJ, SONU AMBWANI and LATA PALIWAL

Biochemical changes associated with triclosan exposure in the freshwater fish *Cyprinus carpio* 140
MADHU SHARMA, YASHPAL, RIJUL RANA, DEEKSHA, DEEPIKA THAKUR and TARANG KUMAR SHAH

Assessment of awareness among respondents towards Pradhan Mantri Jan Dhan Yojana: A study in Haryana 148
MEGHA GOYAL, SUMAN GHALAWAT, MINAKSHI, ATUL DHINGRA and MINI

Youth perception and constraints in agri-entrepreneurship: A factor analytic study 156
JYOTI, SUMAN GHALAWAT, MEGHA GOYAL, NEELAM KAUSHAL, SUNIL KUMAR and VIKRANT HOODA

Design of Object Detection Model Using Deep Learning for Autonomous Driving in Adverse Weather Conditions: A GAN-Based Restoration and Depth-Aware Fusion Framework 165
MUMUKSHU BHATT, SUBODH PRASAD, RAJESH SINGH, ASHOK KUMAR and POOJA TAMTA

A novel mechanical purification approach for *Grewia optiva* Fibre-Reinforced epoxy composites: Effect of fibre geometry on structural performance 172
DEEPA SINGH , NEERAJ BISHT, TARUN KUMAR SAXENA and KAMAL RAWAT

Evaluating Long-Term rainfall and temperature trends across the Shipra River basin 184
POONAM SINGH, MOHAN LAL, DHEERAJ KUMAR, P. K. SINGH and GOVIND VERMA

Rainfall-runoff modelling using soft computing techniques for various watersheds of Madhya Pradesh, India 196
SEEMA KUSHWAHA and PRAVENDRA KUMAR

Prevalence and associated risk factors of functional constipation among the selected adult population of Pantnagar, U.S. Nagar, Uttarakhand, India 204
JYOTI SINGH, NEETU DOBHAL, ARCHANA KUSHWAHA, N.C SHAHI and SABBU SANGEETA

Tracking nutritional transitions: A comparative study of child malnutrition (0–5 years) trends across districts of Madhya Pradesh 212
SHALINI CHAKRABORTY

Determinants influencing tourist selection of rural tourism destinations using Garrett Ranking Technique: Evidence from Supi village, Uttarakhand 221
NEHA PANDEY and KAMAL UPRETI

https://doi.org/10.67063/pjr.vol.24.issue.01.06

Genetic divergence and yield trait impact in French bean (*Phaseolus vulgaris* L.)

BHAWANA MAMGAIN[1*], B.P. NAUTIYAL [2], S.C. PANT[3], VAISHALI THAPA[4], HEIGRUJAM RIYASANA[5] and MAYANK[6]

[1,4,5&6] Institute of Agriculture Training and Research, Dehradun-248001, [2] Basic Science and Humanity, Uttarakhand University of Horticulture and Forestry- 246285, [3] Department of Breeding Uttarakhand University of Horticulture and Forestry- 246285 (Uttarakhand)
[*]Corresponding author's email id: 56682_bhawanamamgai@gbpuat-tec.ac.in

ABSTRACT: The present investigation was undertaken to study Genetic Divergence and Yield Trait Impact in French Bean (*Phaseolus vulgaris* L.). The experiment was conducted at the Vegetable Research and Demonstration Block, Department of Vegetable Science, College of Horticulture, VCSG Uttarakhand University of Horticulture and Forestry, Bharsar, Pauri Garhwal (Uttarakhand). A total of 24 genotypes were evaluated in a randomized block design with three replications. Genotypic correlation coefficients were estimated to determine associations among yield and its component traits, while path coefficient analysis was employed to partition correlations into direct and indirect effects on seed yield per plant. Genetic divergence among genotypes was grouped the genotypes into four distinct clusters, indicating substantial genetic variability. The maximum inter-cluster distance was observed between Cluster I ($D^2 = 2.920$) and Cluster III ($D^2 = 5.571$), reflecting the highest level of genetic divergence. Seed weight, number of primary branches per plant, and seed yield per plot exhibited strong positive direct effects on seed yield per plant and this shows their importance as selection criteria. The identified genetically divergent clusters and key yield-influencing traits can be effectively utilized in future French bean breeding programmes for yield improvement.

Keywords: Components, Correlation analysis, Genetic divergence, French bean, Path coefficient analysis, Yield

French bean (*Phaseolus vulgaris* L.) is one of the most important grain and vegetable legumes cultivated across temperate, subtropical, and tropical regions of the world. It is valued for its tender green pods and dry seeds, which are rich sources of protein, dietary fibre, vitamins, and essential minerals, thereby contributing significantly to nutritional security and farm income (Singh, 2001; Beebe *et al.,* 2013).

Despite its economic importance, the productivity of French bean remains relatively low, largely due to a narrow genetic base and the complex inheritance of yield and its component traits. Seed yield is a polygenic trait governed by several interrelated morphological and physiological characters, whose expression is strongly influenced by environmental conditions (Rana *et al*., 2015). Consequently, direct selection for yield alone is often ineffective, necessitating an understanding of the relationships among yield-contributing traits.

Correlation analysis provides information on the degree and direction of association among characters but does not reveal their true cause effect relationships (Falconer and Mackay, 1996). Path coefficient analysis, originally proposed by Wright (1921) and later refined by Dewey and Lu (1959), addresses this limitation by partitioning correlation coefficients into direct and indirect effects, thereby identifying traits that exert a real and substantial influence on yield. The usefulness of correlation and path analysis in defining effective selection criteria for yield improvement in French bean and related legumes has been well documented (Dutta *et al*., 2018; Santos *et al*., 2017).

Genetic divergence analysis using Mahalanobis D^2 statistics further complements these approaches by assessing genetic diversity and classifying genotypes into distinct clusters to identification of genetically divergent parents is essential for exploiting heterosis and generating broad variability in segregating

populations (Singh, 2001). However, information integrating genetic divergence, character association, and path coefficient analysis in French bean under the mid-hill agro-climatic conditions of Uttarakhand is limited. The unique environmental conditions of this region, characterized by moderate temperatures and variable rainfall, may significantly influence trait expression and interrelationships, making region-specific evaluation essential. Therefore, the present study was undertaken to analyze genetic divergence, correlation, and direct and indirect effects of yield-related traits in French bean genotypes under mid-hill conditions of Uttarakhand, with the objective of identifying key selection traits and genetically diverse parents for future breeding programmes.

MATERIALS AND METHODS

The present investigation was carried out during the kharif season of 2018-2019 at the Vegetable Research and Demonstration Block, Department of Vegetable Science, College of Horticulture, VCSG Uttarakhand University of Horticulture and Forestry, Bharsar, Pauri Garhwal, Uttarakhand. The experimental site represents the mid-hill agroclimatic conditions of Uttarakhand, characterized by moderate temperature and variable rainfall, and is well suited for French bean cultivation.

A total of 24 French bean genotypes, comprising released varieties and germplasm accessions, were evaluated during the cropping season. The experiment was laid out in a Randomized Block Design (RBD) with three replications. The plots of size 3m × 1m and each genotype was grown in individual plots at the spacing 100cm × 50cm. Observations were recorded on growth, phenological, yield, and quality-related traits. The characters studied included days to germination, days to first flowering, days to 50% flowering, days to first harvest, pod maturity duration, plant height (cm), number of primary branches per plant, number of pods per plant, pod length (cm), pod diameter (cm), number of seeds per pod, average pod weight (g), 100-seed weight (g), seed yield per plant (g), seed yield per plot (kg), and protein content (%). Observations were recorded from five randomly selected plants in each plot, and the mean values were used for statistical analysis. Protein content of dry seeds was determined by estimating the nitrogen content as per the modified Kejldhal's method (Jackson, 1965) and multiplying it with the factor 6.25 (Dubetz and Wells, 1968) and expressed on percent basis for each genotype.

Genotypic correlation coefficients were computed to assess the association among different traits following the procedure outlined by Al-Jibouri *et al.* (1958). Path coefficient analysis was performed according to the method suggested by Wright (1921) and elaborated by Dewey and Lu (1959) to partition correlation coefficients into direct and indirect effects on seed yield per plant. Genetic divergence among genotypes was estimated using Mahalanobis D^2 statistics (Mahalanobis, 1936), and clustering of genotypes was carried out using Tocher's method to determine the pattern of genetic diversity. All statistical analyses were performed using SPSS and R software.

RESULTS AND DISSCUSSION

The genetic divergence analysis based on Mahalanobis D^2 statistics grouped the 24 French bean genotypes into four distinct clusters (Table 1), indicating the presence of substantial genetic variability among the evaluated germplasm. Cluster IV accommodated the maximum number of genotypes (8), followed by Cluster III (7) and Cluster II (5), while Cluster I contained the minimum number of genotypes (4). The distribution of genotypes across clusters irrespective of their geographic origin suggests that genetic constitution rather than geographical collection influenced cluster formation, a trend widely reported in French bean diversity studies (Verma *et al*., 2014).

The average intra-cluster distances were lower than the corresponding inter-cluster distances, reflecting greater homogeneity within clusters and pronounced divergence among clusters. Cluster III exhibited the lowest intra-cluster distance (2.598), indicating close genetic similarity among its genotypes, whereas Cluster I recorded the highest intra-cluster distance

(2.920), suggesting comparatively greater variability within that cluster. Similar patterns of intra-cluster variation have been reported in D^2-based diversity analyses of common bean germplasm (Kumar, 2024).

Analysis of inter-cluster distances revealed varying degrees of genetic divergence among clusters. The maximum inter-cluster distance was observed between Cluster I and Cluster III (5.571), followed by Cluster I and Cluster IV (4.970) and Cluster I and Cluster II (4.853), indicating that genotypes belonging to these clusters are genetically most divergent. In contrast, the minimum inter-cluster distance occurred between Cluster II and Cluster III (3.758), reflecting closer genetic affinity between these groups. High inter-cluster distances are particularly important for breeding programmes, as crosses between genetically divergent parents are more likely to generate higher heterosis and wider variability in segregating populations (Chhetri *et al*., 2025).

Overall, the clustering pattern and magnitude of D^2 values confirm the existence of wide genetic divergence among the studied French bean genotypes, highlighting the availability of a broad genetic base that can be effectively exploited for selection and hybridization aimed at yield improvement (Verma *et al.,* 2014).

Mean Performance: The cluster mean analysis revealed substantial variation among the four clusters for all sixteen characters studied, confirming the presence of wide genetic diversity among the French bean genotypes (Fig. 2). Differences in cluster means indicate the relative contribution of component traits to genetic divergence and their potential influence on yield expression, as commonly reported in multivariate diversity studies of grain legumes (Mahalanobis, 1936; Singh, 2001).

Cluster I recorded the lowest mean values for days to germination, days to 50% germination, days to first flowering, and days to 50% flowering, indicating its association with earliness. This cluster also exhibited the highest mean values for number of primary branches per plant, seed weight, protein content, seed yield per plant, and seed yield per plot. The superior yield performance of Cluster I may be attributed to the combined effect of earliness and favourable expression of yield-contributing traits, which has been reported to enhance productivity in French bean and related legumes (Dewey and Lu, 1959; Dutta *et al.,* 2018).

Cluster II showed moderate mean performance for most characters and recorded higher values for plant height, pod length, and number of seeds per pod. However, seed yield per plant and per plot were lower compared to Cluster I, suggesting that individual component traits alone may not directly



Fig. 1: Average intra and inter cluster distance (D^2)

Table 1: Clustering pattern of 24 genotypes of French bean on the basis of genetic divergence

Clusters	No. of genotypes	Genotype name
I	4	Lakshmi, LC-2, Harsil LC-1, LC-1
II	5	IC-049810, IC-199211, EC-755542, IC-199208, EC- 755484
III	7	EC-755510, Solan LC-1, Bean No.-2, IC-84337, EC- 755478, EC-755480, IC-84376
IV	8	755444, EC-755508EC-755509, EC- 755455, EC-755477

translate into higher yield unless supported by other complementary traits. Similar observations have been reported in earlier studies on French bean (Santos *et al.,* 2017).

Cluster III was characterized by relatively later flowering, as indicated by higher mean values for days to first flowering and days to 50% flowering. Although this cluster exhibited a shorter pod set to pod maturity duration, it recorded the lowest seed yield per plant and per plot. The lower yield observed in this cluster may be associated with less favourable combinations of yield-contributing traits rather than any single physiological limitation, and therefore warrants further investigation.

Cluster IV represented late-flowering genotypes and exhibited comparatively higher plant height and moderate average pod weight. Seed yield per plant and per plot were second highest after Cluster I, indicating that delayed phenology does not necessarily limit yield potential when supported by favourable yield attributes. Such genotypes may be useful in breeding programmes aimed at broadening the genetic base and improving yield stability (Singh, 2001; Beebe *et al*., 2013).

Overall, the marked differences in cluster means for yield and related traits suggest that genotypes from genetically divergent clusters, particularly Cluster I in combination with Cluster III or Cluster IV, could be exploited in hybridization programmes to generate wider variability and improved yield potential. However, a more definitive understanding of trait contributions to seed yield requires explicit integration of correlation and path coefficient analyses, including the magnitude of direct and indirect effects of individual traits. The absence of detailed discussion on these relationships represents a limitation of the present study and highlights the need for their comprehensive interpretation alongside cluster mean performance.

CONCLUSION

Based on cluster mean performance, Cluster I emerged as the most promising group, characterized by earliness, higher number of primary branches, greater seed weight, superior protein content, and maximum seed yield per plant and per plot. These traits—particularly seed weight, number of branches per plant, and earliness parameters should be given priority as key selection criteria for yield



Fig. 2: Cluster means for different characters in twenty-four genotypes of French bean

improvement in French bean. Cluster IV, despite its relatively late flowering, also exhibited high yield potential and may serve as a valuable source of favorable yield attributes. Considering the magnitude of inter-cluster divergence and complementary trait expression, genotypes belonging to Cluster I in combination with those from Cluster III or Cluster IV are recommended for hybridization programmes. Such crosses are expected to generate greater heterosis and wider variability in segregating generations, facilitating the development of high-yielding and nutritionally superior French bean varieties. Overall, the findings provide clear trait-based selection guidelines and parent combinations that can be effectively utilized in future French bean breeding efforts.

REFERENCES

Al-Jibouri H. W., Miller, P. A. and Robinson, H. F. (1958). Genotypic and environmental variances and covariances in an upland cotton cross of inter-specific origin. *Agronomy Journal,* 50:633-636.

Beebe, S. E., Rao, I. M., Blair, M. W. and Acosta-Gallegos, J. A. (2013). Phenotyping common beans for adaptation to drought. *Frontiers in Physiology,* 4: 35. https://doi.org/10.3389/fphys.2013.00035

Dubetz, S. and Welis, S. A. (1968). Relation of barley varieties to nitrogen fertilization. *Journal Agriculture Science,* 70:253-256.

Chhetri, M., Sharma, R. and Singh, A. (2025). Genetic divergence and cluster analysis in French bean (*Phaseolus vulgaris* L.) under temperate conditions. *Journal of Plant Genetic Resources, 38*(1): 45–52.

Dewey, D. R. and Lu, K. H. (1959). A correlation and path-coefficient analysis of components of crested wheatgrass seed production. *Agronomy Journal,* 51: 515–518.

Dutta, P., Bera, P. S. and Chakraborty, A. (2018). Genetic diversity analysis in French bean (*Phaseolus vulgaris* L.) using multivariate techniques. *Legume Research,* 41(5): 653–659.

Falconer, D. S. and Mackay, T. F. C. (1996). *Introduction to quantitative genetics* (4th ed.). Longman Group Ltd.

Jackson, M. L. (1965). Soil chemical analysis. Prentice hall of Englewood cliffs, New Jersey, USA.

Kumar, R. (2024). Assessment of genetic diversity in common bean (*Phaseolus vulgaris* L.) using D^2 statistics. *Asian Journal of Plant Breeding,* 9(2): 112–118.

Mahalanobis, P. C. (1936). On the generalized distance in statistics. *Proceedings of the National Institute of Sciences of India,* 2: 49–55.

Rana, J. C., Sharma, T. R., Tyagi, R. K., Chahota, R. K., Gautam, N. K., Singh, M. and Ojha, S. N. (2015). Characterization of Indian common bean germplasm. *Plant Genetic Resources: Characterization and Utilization,* 13(2): 164–174.

Santos, A. D., Ceccon, G., Teodoro, P. E., Correa, A. M. and Silva, J. F. (2017). Genetic diversity and selection of common bean genotypes using multivariate techniques. *Genetics and Molecular Research,* 16(2): gmr16029696. https://doi.org/10.4238/gmr16029696

Singh, S. P. (2001). Broadening the genetic base of common bean cultivars: A review. *Crop Science,* 41(6): 1659–1675. https://doi.org/10.2135/cropsci2001.1659

Upadhyay, D. K., Singh, R. K. and Singh, M. (2019). Genetic divergence studies in French bean (*Phaseolus vulgaris* L.) under mid-hill conditions. *Indian Journal of Agricultural Sciences,* 89(9): 1465–1469.

Verma, A., Singh, B. and Sharma, V. (2014). Genetic divergence studies in French bean (*Phaseolus vulgaris* L.). *Indian Journal of Agricultural Sciences,* 84(7): 829–833.

Wright, S. (1921). Correlation and causation. *Journal of Agricultural Research,* 20: 557–585.

Received: December 27, 2025
Accepted: January 27, 2026